



Effects of fluazuron and ivermectin treatment of cattle on the structure of dung beetle communities

U. Kryger*, C. Deschodt, C.H. Scholtz

Department of Zoology and Entomology, University of Pretoria, 0002 Pretoria, South Africa

Received 27 August 2003; received in revised form 22 July 2004; accepted 4 August 2004

Abstract

A long-term and large-scale field study was carried out to assess the ecotoxicological effect on the dung beetle fauna of cattle treatment with ivermectin (broad-spectrum endectocide) and fluazuron (acaricide) under normal extensive farming conditions in South Africa. One herd of 25 heifers was treated four times at an 8-week interval (starting late November 2001) with a standard injection of ivermectin (200 µg/kg) and a standard pour-on dosage of fluazuron (3 mg/kg) and another herd was left untreated as a control. This set-up was replicated on a second, separate farm and the four herds were kept in four separate paddocks of about 80 ha each. Dung beetle communities were monitored over an entire farming season, from early November 2001 until August 2002. This study was carried out in a very moist year with above average rainfall. The impact of the treatment on the dung beetle communities was examined using a variety of community measures as well as a wide range of univariate, graphical and multivariate analyses. There was no observable effect of the administered drugs on the dung beetle communities. Species richness and diversity seemed unaffected in the treated communities and the ecological similarity of the treated and control communities remained high through most of the trial. These results support the notion that the ecotoxicological impact of antiparasitic drugs depends on factors such as climatic conditions, spatial scale of treatment and proportion of animals treated. © 2004 Elsevier B.V. All rights reserved.

Keywords: Dung beetle communities; Ecotoxicology; Fluazuron; Ivermectin; Long-term field trial

1. Introduction

The majority of South African commercial cattle farmers apply intensive parasite control using various antiparasitic drugs (Spickett and Fivaz, 1992). While

the effective control of parasites seems socio-economically necessary to the farmer, unintended side effects of many pesticides make them environmentally undesirable. Ecotoxicological effects on non-target organisms vary from product to product and scientific experiments aiming to identify products with the least detrimental effects are needed to provide farmers with the necessary information.

Macrocyclic lactones, especially ivermectin, are commonly used because of their broad spectrum against gastro-intestinal nematodes, lungworms, as

* Corresponding author. Tel.: +27 12 420 5343; fax: +27 12 362 5242.

E-mail addresses: ukryger@zoology.up.ac.za (U. Kryger), cdeschodt@zoology.up.ac.za (C. Deschodt), chschooltz@zoology.up.ac.za (C.H. Scholtz).

endectocides with low vertebrate toxicity (Benz et al., 1989; Pulliam and Preston, 1989; Strong and Wall, 1990; Wardhaugh and Beckmann, 1996, 1997).

After administration ivermectin is metabolized only to a limited extent and the parent compound and its metabolites subsequently enter the environment via the treated livestock's faeces (Halley et al., 1989; Wardhaugh and Beckmann, 1996). Concern has been raised about potential adverse effects on dung beetles, which contribute to soil fertility and sustainability of pastoral ecosystems (Wall and Strong, 1987; Strong, 1993; Wardhaugh and Beckmann, 1996, 1997; McKellar, 1997). Laboratory injections of avermectin/ivermectin to heifers negatively affected larval dung beetles including *Diastelopalpus quinquegens* and *Onthophagus gazella* (Sommer et al., 1993; Sommer and Overgaard Nielsen, 1992), *O. binodis* (Roncalli, 1989), *O. ferox* (Ridsdill-Smith, 1993), *Euoniticellus intermedius* (Fincher, 1992) and *Onitis alexis* (Krüger and Scholtz, 1997). Field studies in South Africa detected reduced species diversity and increased species dominance in dung beetle communities after ivermectin treatment of cattle under drought, but not under high-rainfall conditions (Krüger and Scholtz, 1998a,b).

Ivermectin is often used in combination with fluazuron (benzoylphenylurea), a very effective acaricide used to control the blue tick (*Boophilus microplus*) in South Africa. The compound is slowly metabolized and the major route of elimination is via the faeces of treated animals. Bioassays have shown that exposure to fluazuron had no detrimental effect on brood production or egg to adult development of *E. intermedius*, but lead to reduced brood production of *O. gazella*, and *O. taurus* (Fisara, 1994, 1995a,b, 1996).

The objective of this study was to determine the effect of a combined treatment with Acatak[®] (2.5% fluazuron pour-on) and Cevamec[®] (1% ivermectin injectable solution) during an entire grazing season on the structure of the dung beetle community under normal extensive farming conditions in South Africa.

2. Material and methods

This study was conducted on two commercial cattle farms under the same management (Middlepunt and

Abel, 26°54'S, 27°35'E). The farms were 1 km apart 14 km east of Parys (Free State Province, South Africa), at 1350 m.a.s.l. on clay-loam soil. The sites were in the summer rainfall region and the grassland biome of South Africa (Rutherford and Westfall, 1986). Both farms were divided into two adjoining paddocks of 80 ha each with similar vegetation and rainfall patterns; one paddock on each farm being used for the treated cattle and one for the control group. Precipitation and temperature were recorded at the meteorological stations of the South African Weather Bureau at Parys and Vereeniging.

On each farm one herd of 25 cows (with calves) was treated simultaneously with the two products according to the manufacturer's specifications and another herd was left untreated as control group. Acatak[®] (fluazuron 2.5%) was applied as pour-on at 3 mg fluazuron/kg body mass, Cevamec[®] (ivermectin 1%) as subcutaneous injections at 200 µg ivermectin/kg body mass. The treatments were applied in late November 2001, January, March, and May 2002.

Dung beetle populations were sampled on nine occasions in each of the four paddocks along five randomly selected transects with five pitfall traps each. The first sampling occurred just before the first treatment in November 2001, and was repeated at monthly interval until May 2002, once in July and once in August 2002. The trapped beetles were sorted, counted and identified.

2.1. Statistical analyses

The overall diversity between treatment and control paddocks was compared using species richness and Shannon index (Magurran, 1988). The evenness of species abundance was calculated with Pielou's J' ($J' = H'/\ln S$; Pielou, 1975). These indices were computed with the software ESTIMATES (Colwell, 1997).

The distributional properties of each dung beetle community were presented as k -dominance curves (Lambshhead et al., 1983; Magurran, 1988) using the program PRIMER (Plymouth Routines In: Multivariate Ecological Research; Clarke, 1993).

Non-metric multidimensional scaling (MDS) was performed on the fourth-root transformed, mean abundances of species in the communities using the Bray-Curtis coefficient (Bray and Curtis, 1957; Faith et al., 1987) in SAS[®] (SAS Institute Inc.).

Table 1
Measures of overall dung beetle diversity in four South African paddocks

Month	Farm	Regime	<i>S</i>	<i>H'</i>	<i>J'</i>
November-01	Middelpunt	Control	38	2.65	0.72851
	Middelpunt	Cevamec/Acatak	34	2.67	0.75715
	Abel	Control	23	2.63	0.83878
	Abel	Cevamec/Acatak	39	2.25	0.61416
December-01	Middelpunt	Control	32	2.24	0.64633
	Middelpunt	Cevamec/Acatak	33	2.67	0.76362
	Abel	Control	25	2.48	0.77046
	Abel	Cevamec/Acatak	36	2.71	0.75624
January-02	Middelpunt	Control	27	2.33	0.70695
	Middelpunt	Cevamec/Acatak	31	2.05	0.59697
	Abel	Control	35	2.33	0.65535
	Abel	Cevamec/Acatak	35	2.77	0.77911
February-02	Middelpunt	Control	35	2.73	0.76786
	Middelpunt	Cevamec/Acatak	31	2.51	0.73093
	Abel	Control	44	2.75	0.72671
	Abel	Cevamec/Acatak	43	2.76	0.73381
March-02	Middelpunt	Control	31	2.87	0.83576
	Middelpunt	Cevamec/Acatak	35	2.8	0.78755
	Abel	Control	38	2.7	0.74225
	Abel	Cevamec/Acatak	36	2.28	0.63625
April-02	Middelpunt	Control	27	2.03	0.61593
	Middelpunt	Cevamec/Acatak	28	1.94	0.5822
	Abel	Control	30	2.51	0.73798
	Abel	Cevamec/Acatak	29	2.25	0.66819
May-02	Middelpunt	Control	24	2.51	0.78979
	Middelpunt	Cevamec/Acatak	21	2.57	0.84414
	Abel	Control	23	2.63	0.83878
	Abel	Cevamec/Acatak	27	2.03	0.61593
July-02	Middelpunt	Control	8	1.89	0.9089
	Middelpunt	Cevamec/Acatak	10	1.63	0.7079
	Abel	Control	7	1.1	0.56529
	Abel	Cevamec/Acatak	7	1.13	0.58071
August-02	Middelpunt	Control	14	2.34	0.88668
	Middelpunt	Cevamec/Acatak	19	2.44	0.82868
	Abel	Control	15	1.89	0.69792
	Abel	Cevamec/Acatak	15	1.71	0.63145

Species richness, *S*; Shannon's diversity index, *H'*; and species evenness, *J'*. Global *t*-tests showed no significant differences between control and treatment paddocks.

Differences between sampling sites and dates were analyzed for similarity via ANOSIM, a component of PRIMER (Clarke, 1993; Clarke and Gorley, 2001).

The 'similarity percentages' procedure (SIMPER, a component of PRIMER) described by Clarke (1993) and Clarke and Warwick (1994), was used to identify similarity/dissimilarity according to contributions of each species (δ_i) and ranking them in order of importance.

3. Results

A total of 33,388 dung beetle specimens representing 72 species were processed.

During the rain season (October 2001–April 2002), rainfall reached a total of 865.9 mm, clearly above the mean (1904–1994) of 520.5 mm for this period.

The values for the overall diversity of dung beetles shown in Table 1 revealed a relatively stable and




Fig. 1. k -dominance curves for dung beetle abundance in treated and control paddocks of two South African farms on nine sampling dates (a)–(f); (●) = control on farm Abel, (○) = control on farm Middelpunt, (▼) = treated on farm Abel, (▽) = treated on farm Middelpunt.

comparable dung beetle community in both farms. Pre- and post-treatment values for all diversity measures were very homogenous and there was no difference in the dung beetle communities of the control compared to the treatment paddocks. The drop in diversity in May–August 2002 was due to a halt in dung beetle activity during the winter months.

The k -dominance curves were sigmoid for both paddocks on farm Middelpunt in November 2001 (Fig. 1(a)), indicative of undisturbed communities with high species diversity. In December 2001 this situation was similar on farm Abel, with the control paddock showing a lower diversity than the treatment paddock (Fig. 1(b)). This pattern stayed stable for both farms




Fig. 1. (Continued).

until May 2002 (Fig. 1(a)–(c)), suggesting that the dung beetle communities experienced no negative side effects from the treatment of cattle with Acatak[®]/Cevamec[®].

The similarity of the dung beetle communities in both treated and control paddocks in terms of relative species abundance was very high throughout most of the season (Fig. 2). The same level of relative similarity between treatment and control communities as in the pre-treatment sample was found from December 2001 until May 2002, and the communities in the control paddocks were more similar to those of the corresponding treatment than to each other (Fig. 2).

There was a highly significant change in community within each paddock among the various dates. Sampling transects in each paddock were indistinguishable and consequently pooled for further analyses (Table 2). However, in both farms the dung beetle communities before and after treatment were not significantly different from each other, indicative of no negative ecotoxicological effect.

Pooling the data on farm level showed slight indications of differences in the beetle communities between the two farms, six values barely allowing separation, two *R*-values approaching the critical 0.5 limit for clear difference (Table 2). The data were therefore, analyzed separately for each farm (values not shown). The differences in terms of dung beetle communities consistently retrieved significant and high values among sampling dates but control and treatment paddocks remained indistinguishable.

The SIMPER analyses showed that on both farms the number of discriminating species remained within the same range throughout the experiment and did not increase after the treatment. Furthermore, absolute and percentage numbers of typical species were comparable in the control and treated paddocks and remained at the same level in the latter throughout most of the trial. Exceptions to this trend were only found in the winter months on both farms and in March on farm Abel and coincided with the dry period.




Fig. 2. Two-dimensional, non-metric scaling ordination of the relative, fourth-root transformed species abundance data of pooled samples collected on nine different dates; (◇) = control paddocks, (◆) = treated paddocks.

4. Discussion

The data collected over 9-month field study spanning an entire farming season showed that the treatment of cattle with ivermectin and fluzuron had

Table 2
ANOSIM results of testing for differences in dung beetle communities between treatment and control data from two South African farms pooled

	1-way R-value	2-way (Regime/Farm) R-value	2-way (Regime/Date) R-value
Global	–0.001	–0.005/0.039	0.059/ 0.66
November	0.187	0.268/0.37	
December	0.136	0.194/0.188	
January	–0.032	0.06/0.496	
February	0.024	0.144/0.496	
March	–0.02	0.012/0.126	
April	–0.041	–0.04/0.342	
May	0.112	0.13/0.278	
July	0.173	0.146/0.08	
August	–0.006	–0.003/0.276	

Bold = significant differences between communities.

no adverse effect on the dung beetle fauna. Observed disparities between treatment and control were insignificant and probably due to the respective paddocks and their microclimate.

In the present study *O. gazella* did not contribute to differences between treatment and control paddocks, and *E. intermedius* was one of the more common species across all paddocks on both farms. Moreover, *E. intermedius* was abundant in the treated paddocks suggesting that the influence of the pesticides was not marked.

Instead of using the normal application on young animals only, in the present study drugs were administered to all adult animals in the treated herds. As a consequence, the entire dung beetle fauna was exposed synchronously to faecal residues. The absence of any difference in the dung beetle communities between treated and control paddocks strongly suggests that the treatment did not affect the monitored species.

Krüger and Scholtz (1998a,b) found that ivermectin led to changes in dung insect communities under

drought conditions, but not under high-rainfall conditions. The rainfall registered during September 2001 to May 2002 in the present trial was 345.4 mm above average for the rain season. This emphasizes the influence of climatic conditions on the impact of antiparasitic drugs (McKellar, 1997).

The spatial scale of treatment is known to have an important influence on the impact of pesticides on invertebrate populations (Duffield and Aebischer, 1994; Jepson and Thacker, 1990). Plots in this trial were 80 ha, with the two treated plots spatially well separated but each treated plot adjacent to a control plot. Immigration of dung beetles from the control into the treatment paddocks may have moderated any non-lethal effect of the pesticides, but no difference was observed among treated transects.

This study demonstrated that cattle treatment with ivermectin and fluazuron under extensive farming conditions on the South African Highveld can be considered safe with regards to the dung beetle communities under high-rainfall condition.

Acknowledgements

The authors would like to thank Naeem Bham and Barry Hyman (Novartis Animal Health South Africa) for technical advice on product application, Adrian Davis (Department Zoology and Entomology, University Pretoria) for help with dung beetle identifications, and Mike van der Linde (Department of Statistics, University of Pretoria) for the SAS programming.

References

- Benz, G.W., Roncalli, R.A., Gross, S.J., 1989. Use of ivermectin in cattle, sheep, goats and swine. In: Campbell, W.C. (Ed.), *Ivermectin and Abamectin*. Springer, New York, pp. 215–229.
- Bray, J.R., Curtis, J.T., 1957. An ordination of the upland forest of Southern Wisconsin. *Ecol. Monogr.* 27, 325–349.
- Clarke, K.R., 1993. Non-parametric multivariate analyses of changes in community structure. *Aust. J. Ecol.* 18, 117–143.
- Clarke, K.R., Gorley, R.N., 2001. *Primer v5: user manual/tutorial*. PRIMER-E Ltd., Plymouth.
- Clarke, K.R., Warwick, R.M., 1994. Change in Marine Communities: An Approach to Statistical Analysis and Interpretation. Plymouth Marine Laboratory, Plymouth.
- Colwell, R.K., 1997. Estimate S: Statistical estimation of species richness and shared species from samples. Version 5. User's Guide and application at: <http://viceroy.eeb.uconn.edu/estimates>.
- Duffield, S.J., Aebischer, N.J., 1994. The effect of spatial scale of treatment with dimethoate on invertebrate population recovery in winter wheat. *J. Appl. Ecol.* 31, 263–281.
- Faith, D.P., Minchin, P.R., Belbin, L., 1987. Compositional dissimilarity as a robust measure of ecological distance. *Vegetation* 69, 57–68.
- Fincher, G.T., 1992. Injectable ivermectin for cattle: effect on some dung inhabiting insects. *Environ. Entomol.* 21, 871–876.
- Fisara, P., 1994. The effect on dung beetles *Onthophagus taurus*, *Euoniticellus intermedius* and *Onthophagus gazella* of exposure to Fluazuron in cattle faeces. Technical Memorandum No. 94M/10/1466, Ciba-Geigy Australia Ltd., Kemp's Creek.
- Fisara, P., 1995a. The effect on *Onthophagus gazella* exposed to fluazuron in cattle faeces. Technical Memorandum No. 95M/2/1482, Ciba-Geigy Australia Ltd., Kemp's Creek.
- Fisara, P., 1995b. The effects on the dung beetle *Euoniticellus intermedius* of feeding them on faeces derived from cattle treated with ACATAK[®]. Technical Memorandum No. 95/4/1489. Ciba-Geigy Australia Ltd., Kemp's Creek.
- Fisara, P., 1996. The effect on the dung beetle *Euoniticellus intermedius* of feeding them on faeces derived from cattle treated with ACATAK[®] at the rate of 2.5 mg/kg. Technical Memorandum No. 96/6/1528. Ciba-Geigy Australia Ltd., Kemp's Creek.
- Halley, B.A., Nessel, R.J., Lu, A.Y.H., 1989. Environmental aspects of ivermectin usage in livestock: general considerations. In: Campbell, W.C. (Ed.), *Ivermectin and Abamectin*. Springer, New York, pp. 162–172.
- Jepson, P.C., Thacker, J.R.M., 1990. Analysis of the spatial component of pesticide side-effects on non-target invertebrate populations and its relevance to hazard analysis. *Func. Ecol.* 4, 349–355.
- Krüger, K., Scholtz, C.H., 1997. Lethal and sublethal effects of ivermectin on the dung-breeding beetles *Euoniticellus intermedius* (Reiche) and *Onitis alexis* Klug (Coleoptera, Scarabaeidae). *Agric. Ecosyst. Environ.* 61, 123–131.
- Krüger, K., Scholtz, C.H., 1998a. Changes in the structure of dung insect communities after ivermectin usage in a grassland ecosystem. I. Impact of ivermectin under drought conditions. *Acta Oecol.* 19, 425–438.
- Krüger, K., Scholtz, C.H., 1998b. Changes in the structure of dung insect communities after ivermectin usage in a grassland ecosystem. II. Impact of ivermectin under high-rainfall conditions. *Acta Oecol.* 19, 439–451.
- Lambhead, P.J.D., Platt, H.M., Shaw, K.M., 1983. The detection of differences among assemblages of marine benthic species based on an assessment of dominance and diversity. *J. Nat. Hist.* 17, 859–874.
- Magurran, A.E., 1988. *Ecological Diversity and its Measurements*. Groom Helm, London.
- McKellar, Q.A., 1997. Ecotoxicology and residues of anthelmintic compounds. *Vet. Parasitol.* 72, 413–435.
- Pielou, E.C., 1975. *Ecological Diversity*. Wiley, New York.

- Pulliam, J.D., Preston, J.M., 1989. Safety of ivermectin in target animals. In: Campbell, W.C. (Ed.), *Ivermectin and Abamectin*. Springer, New York, pp. 149–161.
- Ridsdill-Smith, T.J., 1993. Effects of avermectin residues in cattle dung on dung beetle (Coleoptera: Scarabaeidae) reproduction and survival. *Vet. Parasitol.* 48, 127–137.
- Roncalli, R.A., 1989. Environmental aspects of use of ivermectin and abamectin in livestock: effects on cattle dung fauna. In: Campbell, W.C. (Ed.), *Ivermectin and Abamectin*. Springer, New York, pp. 173–181.
- Rutherford, M.C., Westfall, R.H., 1986. Biomes of southern Africa—an objective categorization. *Mem. Bot. Surv. S. Afr.* No. 54.
- Sommer, C., Gronvold, J., Holter, P., Nansen, P., 1993. Effects of ivermectin on two afro-tropical dung beetles, *Onthophagus gazella* and *Diastelopalpus quinquedens* (Coleoptera: Scarabaeidae). *Vet. Parasitol.* 48, 171–179.
- Sommer, C., Overgaard Nielsen, B., 1992. Larvae of the dung beetle *Onthophagus gazella* Fabricius (Coleoptera: Scarabaeidae) exposed to lethal and sublethal ivermectin concentrations. *J. Appl. Entomol.* 114, 502–509.
- Spickett, A.M., Fivaz, B.H., 1992. A survey of cattle tick control practices in the Eastern Cape province of South Africa. *Onderstepoort J. Vet. Res.* 59, 203–210.
- Strong, L., 1993. Overview: the impact of avermectins on pasture-land ecology. *Vet. Parasitol.* 48, 3–17.
- Strong, L., Wall, R., 1990. The chemical control of livestock parasites: problems and alternatives. *Parasitol. Today* 9, 291–296.
- Wall, R., Strong, L., 1987. Environmental consequences of treating cattle with the antiparasitic drug ivermectin. *Nature* 324, 418–420.
- Wardhaugh, K., Beckmann, R., 1996. Chemicals affect soil animals. *Rural Res.* 173, 7–10.
- Wardhaugh, K., Beckmann, R., 1997. Chemicals affect soil animals. *Rural Res.* 173, 7–10.